

PEACH

- ✓ Absolute lack of latency periods
- ✓ Fruits consistent
- ✓ Post-harvest long life of fruits
- ✓ Improving the coloring of the fruit
- ✓ Increased plant endogenous resistance to stress
- ✓ Vegetative balance
- ✓ Increase in Brix grades
- ✓ Homogeneous flowering
- ✓ Greater resistance to diseases

| RADICAL | N. APPLICATIONS | DOSAGE |
|---------|---|--|
| DUST | 1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)* | 400/475 Kg per Ha 400/475 Kg per Ha |
| COMPLEX | 1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)* | 800/1000 Kg per Ha 800/1000 Kg per Ha |
| EVO | 1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)* | 750/950 Kg per Ha 750/950 Kg per Ha |
| PRO | 1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)* | 650/825 Kg per Ha 650/825 Kg per Ha |

Locate near the roots of the plants - bury the product after distribution

| FOLIAR | N. APPLICATIONS | DOSAGE |
|-----------|--|---|
| SUPERFINE | PRE-FLOWERING (3 APPLICATIONS) DURING FLOWERING (1-2 APPLICATIONS) POST FLOWERING (5-8 APPLICATIONS) | 5 Kg per Ha (a app.) 5 Kg per Ha (a app.) 3 Kg per Ha (a app.) |

* Do not use chemical fertilizers in the soil where Abomin is applied to the soil, they would counteract its effectiveness.