



APPLE

- ✓ Improved productivity and grade
- ✓ Summer pruning reduced or completely removed
- ✓ Greater orchard vegetation-production balance
- ✓ Plant growth less stressed
- ✓ Increased colouring on red apples
- ✓ Greater resistance to disease
- ✓ Increased sugar content (Brix)

RADICAL	N. APPLICATIONS	DOSAGE
GRANULAR	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	375 Kg per Ha 375 Kg per Ha
COMPLEX	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	800 Kg per Ha 800 Kg per Ha
PRO	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	650 Kg per Ha 650 Kg per Ha

Locate near the roots of the plants - bury the product after distribution

FOLIAR	N. APPLICATIONS	DOSAGE
ULTRAFINE	PRE-FLOWERING (3 APPLICATIONS) DURING FLOWERING (1-2 APPLICATIONS) POST FLOWERING (5-8 APPLICATIONS)	5 Kg per Ha (a app.) 5 Kg per Ha (a app.) 3 Kg per Ha (a app.)

* Do not use chemical fertilizers in the soil where Abomin is applied to the soil, they would counteract its effectiveness.

