

- ✓ Vegetative balance
- Correction of any deficiencies
- Homogeneous flowering
- Firm fruit (high specific weight)
- ✓ Long life after fruit harvest

- Greater resistance to various stresses
- Improvement of the colours of the fruit
- Higher yield per hectare higher productivity
- Greater size of the fruits and their homogenization
- Greater resistance to disease

RADICAL	N. APPLICATIONS	DOSAGE
GRANULAR	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	325 Kg per Ha 325 Kg per Ha
COMPLEX	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	700 Kg per Ha 700 Kg per Ha
PRO	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	575 Kg per Ha 575 Kg per Ha

Locate near the roots of the plants - bury the product after distribution

FOLIAR	N. APPLICATIONS	DOSAGE
ULTRAFINE	PRE-FLOWERING (3 APPLICATIONS) DURING FLOWERING (1-2 APPLICATIONS) POST FLOWERING (5-8 APPLICATIONS)	5 Kg per Ha (a app.)5 Kg per Ha (a app.)3 Kg per Ha (a app.)

^{*} Do not use chemical fertilizers in the soil where Abomin is applied to the soil, they would counteract its effectiveness.



