

# CITRUS FRUIT

- ✓ Productivity increased
- ✓ Early maturation and increase in Brix grades
- ✓ Absolute lack of latency periods
- ✓ Vegetative balance
- ✓ More intense coloring of the fruits
- ✓ Increase of the shelf-life of the fruits also in the cold room
- ✓ More homogeneous flowering
- ✓ Greater resistance to diseases

RADICAL	N. APPLICATIONS	DOSAGE
GRANULAR	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	<b>375</b> Kg per Ha <b>375</b> Kg per Ha
COMPLEX	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	<b>800</b> Kg per Ha <b>800</b> Kg per Ha
PRO	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	<b>650</b> Kg per Ha <b>650</b> Kg per Ha

Locate near the roots of the plants - bury the product after distribution

FOLIAR	N. APPLICATIONS	DOSAGE
ULTRAFINE	PRE-FLOWERING (3 APPLICATIONS) DURING FLOWERING (1-2 APPLICATIONS) POST FLOWERING (5-8 APPLICATIONS)	<b>5</b> Kg per Ha (a app.) <b>5</b> Kg per Ha (a app.) <b>3</b> Kg per Ha (a app.)

\* Do not use chemical fertilizers in the soil where Abomin is applied to the soil, they would counteract its effectiveness.

