

TABLE GRAPES

- ✔ Productivity increased
- ✔ Increase of the berry polar and equatorial diameters
- ✔ Higher bunch average weight and grape homogeneity
- ✔ Less manual work of grape selection (grape-stones)
- ✔ Early maturation and increase in Brix grades
- ✔ Increased plant endogenous resistance to thermal and biotic stress
- ✔ Improved and early lignification of shoots
- ✔ Improved chance of selecting wood to prune
- ✔ Greater vineyard vegetation-production balance

RADICAL	N. APPLICATIONS	DOSAGE
GRANULAR	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	400 Kg per Ha 400 Kg per Ha
COMPLEX	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	800 Kg per Ha 800 Kg per Ha
PRO	1° APPLICATION (autumnal)* 2° APPLICATION (pre-flowering)*	650 Kg per Ha 650 Kg per Ha

Locate near the roots of the plants - bury the product after distribution

FOLIAR	N. APPLICATIONS	DOSAGE
ULTRAFINE	PRE-FLOWERING (3 APPLICATIONS) DURING FLOWERING (1-2 APPLICATIONS) POST FLOWERING (5-8 APPLICATIONS)	5 Kg per Ha (a app.) 5 Kg per Ha (a app.) 3 Kg per Ha (a app.)

* Do not use chemical fertilizers in the soil where Abomin is applied to the soil, they would counteract its effectiveness.

