

# KIWI

- ✓ Increase in % dry matter
- ✓ Vegetative balance
- ✓ Absolute absence of deficiencies
- ✓ Healthier leaf
- ✓ No alternation in productions
- ✓ Greater vegetative-productive balance of the orchard
- ✓ Greater resistance to diseases

RADICAL	N. APPLICATIONS	DOSAGE
GRANULAR	1° APPLICATION (autumnal)* 2° APPLICATION (Pre-flowering)*	<b>500</b> Kg per Ha <b>500</b> Kg per Ha
COMPLEX	1° APPLICATION (autumnal)* 2° APPLICATION (Pre-flowering)*	<b>1000</b> Kg per Ha <b>1000</b> Kg per Ha
PRO	1° APPLICATION (autumnal)* 2° APPLICATION (Pre-flowering)*	<b>825</b> Kg per Ha <b>825</b> Kg per Ha

Locate near the roots of the plants - bury the product after distribution

FOLIAR	N. APPLICATIONS	DOSAGE
ULTRAFINE	PRE-FLOWERING (3 APPLICATIONS) DURING FLOWERING (1-2 APPLICATIONS) POST FLOWERING (5-8 APPLICATIONS)	<b>5</b> Kg per Ha (ad app.) <b>5</b> Kg per Ha (ad app.) <b>3</b> Kg per Ha (ad app.)

\* Do not use chemical fertilizers in the soil where Abomin is applied to the soil, they would counteract its effectiveness.

